



2560 - Unimodal Palindromic Decompositions

North America - Greater New York - 2002/2003

A sequence of positive integers is *Palindromic* if it reads the same forward and backward. For example:

23 11 15 1 37 37 1 15 11 23

1 1 2 3 4 7 7 10 7 7 4 3 2 1 1

A *Palindromic* sequence is *Unimodal Palindromic* if the values do not decrease up to the middle value and then (since the sequence is palindromic) do not increase from the middle to the end. For example, the first example sequence above is **NOT** *Unimodal Palindromic* while the second example is.

A *Unimodal Palindromic* sequence is a *Unimodal Palindromic Decomposition* of an integer N , if the sum of the integers in the sequence is N . For example, all of the *Unimodal Palindromic Decompositions* of the first few integers are given below:

1: (1)
2: (2), (1 1)
3: (3), (1 1 1)
4: (4), (1 2 1), (2 2), (1 1 1 1)
5: (5), (1 3 1), (1 1 1 1 1)
6: (6), (1 4 1), (2 2 2), (1 1 2 1 1), (3 3),
(1 2 2 1), (1 1 1 1 1 1)
7: (7), (1 5 1), (2 3 2), (1 1 3 1 1), (1 1 1 1 1 1 1)
8: (8), (1 6 1), (2 4 2), (1 1 4 1 1), (1 2 2 2 1),
(1 1 1 2 1 1 1), (4 4), (1 3 3 1), (2 2 2 2),
(1 1 2 2 1 1), (1 1 1 1 1 1 1 1)

Write a program, which computes the number of *Unimodal Palindromic Decompositions* of an integer.

Input

Input consists of a sequence of positive integers, one per line ending with a '0' (zero) indicating the end.

Output

For each input value except the last, the output is a line containing the input value followed by a space, then the number of *Unimodal Palindromic Decompositions* of the input value.

Sample Input

2
3
4
5
6
7
8
10
23
24
131

213
92
0

Sample Output

2 2
3 2
4 4
5 3
6 7
7 5
8 11
10 17
23 104
24 199
131 5010688
213 1055852590
92 331143

Greater New York 2002-2003